

This datasheet offers general information. Please contact your local ECO representative for country-specific details.



OXYTET FG200

Presentation

Oxytet FG200 provides 200g oxytetracycline hydrochloride activity per kg, as a feed stable dihydrate salt.

Indications

Poultry (including chickens, turkeys, ducks)

Treatment and control of a wide variety of bacterial diseases caused by organisms sensitive to oxytetracycline, including: Colisepticaemia, Duck Septicaemia, Omphalitis, Fowl Cholera, Enteritis (*E. coli*, *Clostridium* spp, *Salmonella* spp), Infectious Coryza (*Haemophilus paragallinarum*) and Chronic Respiratory Disease (*Mycoplasma gallisepticum*).

Oxytet FG200 is also useful in the treatment and control of diseases caused by susceptible secondary infections after primary viral, coccidial (*Eimeria* spp) or mycoplasmal infections, at times of stress and when non-specific early mortality is a problem.

Pigs

Treatment and control of respiratory and enteric bacterial diseases caused by oxytetracycline- sensitive organisms, such as: atrophic rhinitis, enzootic pneumonia, *Haemophilus pneumonia*, necrotic enteritis, Porcine Proliferative Enteropathy/ileitis and diseases associated with *E. coli* and *Salmonella choleraesuis*.

Oxytet FG200 is also useful in the treatment and control of diseases due to susceptible secondary infections after a primary viral or mycoplasmal infection.

Calves, lambs and kids

For the control and treatment of respiratory, septicaemic and enteric diseases caused by oxytetracycline-sensitive organisms including those caused by *Salmonella* spp and *Pasteurella* spp and diseases caused by susceptible secondary infections after a primary viral or mycoplasma infection.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Dosage and administration

To achieve therapeutic levels of oxytetracycline, Oxytet FG200 should be mixed in the feed at the following inclusion rate:

Species	Oxytetracycline g per tonne (ppm)	Quantity of Oxytet FG200 kg per tonne of feed
Poultry	100 – 200 g	0.5 – 1.0 kg
Pigs	100 – 400 g	0.5 – 2.0 kg
Lambs/calves/kids	50 – 150 g	0.25 – 0.75 kg

These are average doses suitable for routine use. In the event of severe disease challenge, or when disease causes inappetency, levels can be safely increased to give up to 1000mg oxytetracycline per kg of feed. Treatment should be continued for a period of 5 – 14 days, as directed by a veterinary surgeon.

Accurate mixing is essential. To ensure uniform dispersion, mix the required amount of Oxytet FG 200 with 10 times its weight of feed material before blending into the final feed. During treatment only oxytetracycline-medicated feed should be used.

Contra-indications/warnings etc.

During treatment only oxytetracycline-medicated feed should be used.

Withdrawal period

Animals should not be slaughtered for human consumption during treatment or for 5 days following the end of treatment.

Operator warnings

Avoid direct contact of Oxytet FG200 with the skin or eye. In case of eye contact, immediately flush out with plenty of water. Wear overalls, dust mask, gloves and goggles when handling product. Wash hands and exposed skin after working and before meals.

Pharmaceutical precautions

Store in a cool, dry place, protected from direct sunlight. Ensure bag is sealed securely after each use.

General precautions

For animal treatment only. Keep out of reach and sight of children.

Package quantities

Oxytet FG200 is packed in 25kg bags.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Container disposal

Dispose of waste and empty packaging safely and in accordance with local waste disposal regulations.

Further information

ECO Animal Health
PO Box 47542
London N14 6WS
United Kingdom

Tel: +44 (0)208 447 8899
Fax: +44 (0)208 447 9292
Website: www.ecoanimalhealth.com