

FLAVECO 80

Presentation

Flaveco 80 is a premix for incorporation in feedingstuffs containing 80g flavophospholipol per kg.

Flavophospholipol has several synonyms including bambermycin, moenomycin and flavomycin.

Indications

To improve the rate of growth and efficiency of feed conversion in pigs, poultry, calves, fattening cattle, animals bred for fur and rabbits.

Dosage and administration

For oral administration in feed only.

Stock/Type of feed	Maximum age	Rate of inclusion of Flaveco 80 per tonne of feed	Amount of flavophospholipol per tonne of final feed
Poultry Laying hens Turkey Other poultry: (breeders, broilers, etc) <i>excluding ducks, geese and pigeons</i>	---- 26 weeks 16 weeks	25 – 62.5g 12.5 – 250g 12.5 – 250g	2 – 5g 1 – 20g 1 – 20g
Pigs Milk replacer Dry feed	3 months 6 months	125 – 312g 12.5 – 250g	10 – 25g 1 – 20g
Calves Milk replacer Dry feed	6 months 6 months	100 – 200g 75 – 200g	8– 16g 6 – 16g
Fattening Cattle Concentrate	---	25 – 125g	2 – 10g based on complete feeding stuff
Animals bred for fur	---	25 – 50g	2 – 4g
Rabbits	---	25 – 50g	2 – 4g

Feed block: Add 1kg of the product to 1 tonne of feed to provide 80 mg/kg (80 ppm) flavophospholipol in the finished feed.

As with all feed blocks formulations there is a degree of variation in intake by individual animals.

For supplementary feeding the maximum dose in the daily ration must not exceed the following: 40 mg flavophospholipol for 100 kg bodyweight. Above 100 kg, add 1.5 mg flavophospholipol for each additional 10 kg of bodyweight.

To allow even distribution in the final feed, Flaveco 80 should be mixed in a small quantity of feed prior to incorporating in the final mix.

Feed continuously.

Contra-indications, warnings etc.

Withdrawal period

Bambermycin is not absorbed from the gastro-intestinal tract and therefore no withdrawal period is necessary when fed Flaveco 80.

Operator warnings

Avoid direct contact with skin.

Contra Indications

Not to be used for ducks, geese and pigeons.

Not to be used in breeding stock or adult ruminants other than fattening cattle.

Do not mix or use simultaneously with any feedingstuff containing any other antibiotic or growth promoter.

General precautions

If this product is incorporated in an intermediate feedingstuff, care should be taken to ensure that the intermediate feedingstuff is incorporated at a rate which will yield the same concentration of active ingredient in the final feed as those in the dosage schedule.

As with all feedblock formulations, there is a degree of variation in intake by individual animals.

For animal treatment only. Keep out of reach and sight of children

Storage and stability

Store in a cool, dry place below 25°C. Close bag after use.

The product will remain stable in the finished feed for three months.

The product can be incorporated into pelleted feed preconditioned with steam for up to 5 minutes at a temperature not greater than 70°C .

Package quantities

Flaveco 80 is presented in 20 kg polyethylene-lined, woven polypropylene export quality sacks which are sealed by stitching over tape.

Container disposal

Do not leave or dispose of Flaveco 80 or feed containing Flaveco 80 where it may be accessible to animals not under treatment, or wildlife.

Further information

For further information contact:

ECO Animal Health
PO Box 47542
London N14 6WS
United Kingdom

Tel: +44 (0)208 447 8899
Fax: +44 (0)208 447 9292
Website: www.ecoanimalhealth.com