



SALECO 120

Presentation

Saleco 120 contains 120g salinomycin sodium per kg, for incorporation into animal feedingstuffs.

Indications

Poultry

As an aid in the prevention of coccidiosis in broiler chickens caused by *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. mivati*, *E. necatrix* and *E. tenella*.

Pigs

For use in growing and fattening pigs for the improvement of liveweight gain and feed conversion efficiency. Also, to improve crude protein digestibility and daily nitrogen retention.

Dosage and administration

For administration in feed only.

Broiler chickens

500 g Saleco 120 to be thoroughly mixed into each 1,000 kg of broiler feed to provide 60 mg salinomycin sodium per kg in finished feed (60 ppm).

Pigs

For piglets up to 30 – 40 kg bodyweight (max four months of age). To be incorporated in the feed at a rate of 250-500 g Saleco 120 per tonne to provide 30-60 mg salinomycin sodium per kg (30-60 ppm).

For pigs up to slaughter weight (max. to six months of age): To be incorporated in the feed at a rate of 125-250 g Saleco 120 per tonne to provide 15-30 mg salinomycin sodium per kg (15-30ppm).

To allow thorough dispersion of Saleco 120, it should be mixed with a suitable quantity of feed before incorporation into the final mix.

Contra-indications, warnings etc

Withdrawal period

Birds should not be slaughtered for human consumption during treatment or for 5 days thereafter.

There is no withdrawal period for pigs.

Operator warnings

Operators handling Saleco 120 should wear face masks, gloves and protective overalls.

Avoid inhalation and direct skin contact. Wash thoroughly after handling. If accidental eye contact occurs, immediately rinse with water.

Pharmaceutical precautions

Accurate mixing is essential.

Not for direct feeding.

This coccidiostat must not be mixed or used simultaneously with any other anti-coccidial.

Animals, including birds should not be treated with products containing tiamulin during or for 7 days before or after receiving feed containing Saleco 120. Severe growth depression or death may occur.

Do not allow horses or other equines access to feed containing Saleco 120.

Ingestion of salinomycin by horses has been fatal.

Do not feed to turkeys.

Storage

Store below 25°C in a dry place.

Close container after use.

General precautions

For animal treatment only.

Keep out of reach and sight of children.

Packaging quantities

Net weight 25 kg.

Container disposal

Do not leave or dispose of Saleco 120 or feed containing it where it may be accessible to animals not under treatment or to wildlife.

Further information

For further information contact:

ECO Animal Health
PO Box 47542
London N14 6WS
United Kingdom

Tel: +44 (0)208 447 8899

Fax: +44 (0)208 447 9292

Website: www.ecoanimalhealth.com