

This datasheet offers general information. Please contact your local ECO representative for country-specific details.



ECOTRAZ 20 Pig Pour-On

Presentation

A liquid pour-on for pigs containing 2.0% w/v amitraz.

Indications

Ecotraz 20 Pig Pour-On is used for the control of mange and lice infestations on pigs.

Dosage and administration

Ecotraz 20 Pig Pour-On is a topical pour-on formulation for external use only. The dose is 1ml per 4kg liveweight.

Pig	Dorsal/midline dose	Ears	Total dose
Weaners (10-25kg)	10 ml	2.5 ml/ear	15 ml
Growers (25-60kg 60-100kg)	15 ml	2.5 ml/ear	20 ml
	25 ml	2.5 ml/ear	30 ml
Gilt/small sow (100-180kg)	40 ml	5 ml/ear	50 ml
Sow/boar (180-240kg)	60 ml	5 ml/ear	70 ml

The required dose is applied evenly along the backline from the ears to the tail. A small amount of the calculated dose should be used in each ear to prevent re-infestation.

There is no need to remove any scabs or wax prior to treatment with Ecotraz 20 Pig Pour-on. Can be used simultaneously with anthelmintics and other medications. Shake well before use.

Mange:

For routine treatment of mange, a single treatment is recommended. Where no routine treatment has been made for some time, and in severe cases of mange, apply 2 treatments 7-10 days apart. Where extensive lesions are present, for example ears or legs, spot treat with extra 5ml doses.

Lice:

For complete control of lice, apply 2 treatments 10-14 days apart.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Recommended routine control programme:

- **Sows and Gilts** - treat 2-3 days before farrowing to ensure maximum protection of the piglets. The sow is the main source of mange mites on piglets. Treating sows effectively before farrowing stops the transfer of mange to the piglets resulting in a healthier litter.
- **Boars** - treat every 3 months. Regular treatment of boars will prevent re-infection of sows at mating.
- **Young pigs** - treat after weaning and before moving to fattening houses.
- **New additions to the herd** - isolate new pigs and treat before introducing them into the herd. A second treatment 7-10 days later may be needed.

Infested animal housing should be treated with a suitable amitraz acaricide spray.

Contra-indications, warnings etc

Withdrawal periods

- Pigs treated with Ecotraz 20 Pig Pour-on should not be slaughtered for human consumption until at least 7 days after the last treatment.

Operator warnings

- Wear overalls, rubber gloves and face shield when opening containers or diluting, mixing or transferring the product from one container to another or when washing out the container.
- Wear arm-length rubber gloves, rubber boots and bib apron when handling diluted fluid or freshly treated animals.
- When using do not eat, drink or smoke.
- Do not breathe spray mist.
- Avoid all contact with mouth, skin and eyes.
- Wash concentrate or splashes of diluted spray from skin or eyes immediately.
- Wash all protective clothing thoroughly after use especially the inside of gloves.
- Wash hands and exposed skin before eating, drinking or smoking and after work.
- If you feel unwell seek medical advice and show this document.

Guide to Doctor

- Amitraz is **NOT** an organophosphate compound. If poisoning is suspected treatment should be symptomatic and supportive, paying particular attention to monitoring of respiratory and cardiac function.
- Do **NOT** use Atropine.
- Do **NOT** induce vomiting.

Protection of wildlife and the environment

Harmful to fish. Do not contaminate ponds, waterways or ditches with chemical or used container.

Hazard warnings

May cause sensitisation by skin contact. Harmful if swallowed.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Pharmaceutical precautions

- Close container after use.
- Store below 25°C in a dry place and protect from freezing.
- Keep away from food, drink and animal feedingstuffs.
- Keep out of the reach and sight of children.
- For animal treatment only.
- For external use only.
- Keep away from domestic animals.

General precautions

- Do not use on horses, dogs and cats.

Package quantities

1 litre containers.

Further information

For further information contact:

ECO Animal Health
PO Box 47542
London N14 6WS
United Kingdom

Tel: +44 (0)208 447 8899
Fax: +44 (0)208 447 9292
Website: www.ecoanimalhealth.com