

This datasheet offers general information. Please contact your local ECO representative for country-specific details.



ECOMINTIC 100

Presentation

An aqueous suspension for oral administration of 100mg fenbendazole per ml.

Indications

A broad spectrum oral anthelmintic with ovicidal, larvicidal and adulticidal action for use in cattle, sheep, goats and horses.

Kills worm eggs present in the animal at the time of dosing.
Class I: 100% effective in more than 80% of treated animals

The list contains the more important endoparasite species. This anthelmintic may also be effective against other species. For more information consult your veterinarian.

Worm species	Immature	Adult
Cattle		
<i>Haemonchus</i> spp	*	*
<i>Ostertagia</i> spp	* *	*
<i>Oesophagostomum</i> spp	*	*
<i>Bunostomum phlebotomum</i>	*	*
<i>Dictyocaulus</i> spp	*	*
<i>Cooperia</i> spp	*	*
Sheep and goats		
<i>Haemonchus</i> spp	*	*
<i>Ostertagia</i> spp	*	*
<i>Trichostrongylus</i> spp	*	*
<i>Nematodirus spathiger</i>	*	*
<i>Strongyloides papillosus</i>	*	*
<i>Gaigeria pachyscelis</i>	*	*
<i>Chabertia ovina</i>	*	*
<i>Oesophagostomum</i> spp	*	*
<i>Dictyocaulus</i> spp	*	*
<i>Moniezia expansa</i>	*	*

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Horses		
<i>Strongylus edentatus</i>	*	*
<i>S. equinus</i>	*	*
<i>S. vulgaris</i>	*	*
<i>Triodontophorus</i> spp	*	*
<i>Cyathostomum</i> spp	*	*
<i>Cylicocyclus</i> spp	*	*
<i>Cylicostephanus</i> spp	*	*
<i>Cylicodontophorus</i> spp	*	*
<i>Oxyuris equi</i>	*	*
<i>Parascaris equorum</i>	**	**

* Control = ≥ 90% effective ** 10mg/kg

Dosage and administration

Administer by the oral route as a drench using a syringe or dosing gun. Read label directions carefully before using the product. Use only as directed. For veterinary use only. Shake well before use.

Cattle and calves: 1 ml / 20 kg bodyweight = 5 mg/kg
 Type II ostertagiasis:..... 1 ml / 10 kg bodyweight = 10mg/kg
Sheep and goats: 1 ml / 20 kg bodyweight = 5 mg/kg
Horses: 1 ml / 20 kg bodyweight = 5 mg/kg
 Horse ascarids: 1 ml / 10 kg bodyweight = 10mg/kg

Contra-indications, warnings etc

Withdrawal periods

- **Sheep and goats:** do not slaughter animals for human consumption within **7 days** of treatment.
- **Cattle:** do not slaughter animals for human consumption within **14 days** of treatment.
- **Dairy cattle:** do not consume milk within **3 days** of treatment.
- Do not use in horses intended for human consumption

Operator warnings

- Direct contact with skin should be kept to a minimum.
- Wear suitable protective clothing including impermeable rubber gloves.
- Wash hands after use.
- In case of suspected intoxication, seek medical attention showing the doctor the label.

Protection of wildlife and the environment

- Do not contaminate ponds, waterways or ditches with product or used containers.
- Dispose of used containers safely.
- Wash out containers.
- Do not re-use containers.
- Dispose of surplus product or waste material in accordance with local regulations.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Hazard warnings

May cause sensitisation by skin contact. Harmful if swallowed.

Pharmaceutical precautions

- Close container after use.
- Store below 25°C in a dry place and protect from light.
- Do not freeze.
- Keep away from food, drink, domestic animals and domestic habitation.
- Shake container before use
- For animal treatment only.
- Keep out of the reach and sight of children.

General precautions

- Assess bodyweight as accurately as possible before calculating the dosage.
- Intensive use or misuse of anthelmintics can give rise to resistance. To reduce this risk dosing programmes should be discussed with your veterinary adviser.

Package quantities

1 litre and 5 litre containers.

Further information

For further information contact:

ECO Animal Health
PO Box 47542
London N14 6WS
United Kingdom

Tel: +44 (0)208 447 8899
Fax: +44 (0)208 447 9292
Website: www.ecoanimalhealth.com