

This datasheet offers general information. Please contact your local ECO representative for country-specific details.



AIVLOSIN FG10 (also known as Aivlosin 8.5 mg/g Premix)

Presentation

Aivlosin[®] FG10 (8.5 mg/g) is a free-flowing microgranulated premix for the medication of feedingstuffs, and contains tylvalosin (as tylvalosin tartrate). This provides 10 mg/g of activity when measured by microbiological assay or 8.5mg/g by HPLC analysis.

Properties

Tylvalosin is a new macrolide antibiotic based on a highly active lactone ring. It has antibacterial activity against Gram-positive and some Gram-negative organisms and against mycoplasma by way of inhibition of protein synthesis in the bacteria cell. It is highly effective against a range of important diseases in pigs and poultry

Tylvalosin has both bacteriostatic and bactericidal action plus the additional benefit of antimicrobial activity of the major metabolite 3-AT.

Tylvalosin can act intracellularly and reaches relatively high intracellular concentrations in phagocytic cells (macrophages and neutrophils). The activity of macrophages can be specifically increased and this may help the innate (non-specific) immune system remove foreign material such as pathogens.

Indications

Swine

Aivlosin[®] FG10 (8.5 mg/g) is indicated for the treatment and prevention of Swine Enzootic Pneumonia caused by *Mycoplasma hyopneumoniae*, Ileitis (Porcine Proliferative Enteropathy) caused by the intracellular bacterium *Lawsonia intracellularis* and Swine Dysentery caused by *Brachyspira hyodysenteriae*. Aivlosin[®] is also active against *Brachyspira pilosicoli*, the cause of colitis in pigs.

Poultry

Aivlosin[®] FG10 (8.5 mg/g) is indicated for prevention and treatment of Mycoplasmosis (*Mycoplasma gallisepticum*, *M. synoviae* and other *Mycoplasma* species), and for diseases associated with *Clostridium perfringens* (enteritis resulting in wet litter syndrome and cholangiohepatitis) in chickens, replacement pullets and turkeys. Aivlosin[®] also has activity against *Ornithobacterium rhinotracheale* (ORT) of poultry.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Dosage and administration

Aivlosin FG10 (8.5 mg/g) should be mixed in the feed at the following dose rates.

	Amount of active per ton of feed	Amount of product per ton of feed
Pigs and Poultry – Prevention	50 g (50 ppm)	5 kg
Pigs and Poultry – Treatment	50-100 g (50-100 ppm)	5 – 10 kg

* Consultation with your Veterinary Surgeon is recommended before commencing treatment.

Synergy of tylvalosin and tetracyclines has been shown and programs in the field combining tetracyclines and Aivlosin at lower doses have proven effective.

Swine

For prevention, feed medicated feeding stuff at 50ppm for 21 days or until the end of the risk period.

For treatment, feed medicated feeding stuff at 50 - 100ppm for 10 days. Repeat as necessary.

For pigs with reduced or restricted food intake, inclusion levels may need to be increased to achieve target dosage.

Poultry

For use in a control programme, feed medicated feeding stuff at 50-100ppm for the first 7 days of life, followed by Aivlosin FG10 at these levels for a further 5-7 days during periods of stress (e.g. vaccination).

Contra-indications, warnings etc.

Withdrawal period

Pigs: 2 days

Chickens: 3 days. Not intended for hens that produce eggs for human consumption.

Operator warnings

When mixing the veterinary medicinal product and handling the medicated feed, direct contact with eyes, skin and mucous membranes should be avoided. Wash hands and exposed skin thoroughly after use. In case of eye contact, immediately flush out with plenty of water. In case of accidental ingestion seek immediate medical advice and show the label to the physician. People with a known hypersensitivity to tylvalosin tartrate should avoid any contact with the product.

Pharmaceutical precautions

Do not store product above 25°C. Store in a cool, dry place, protected from direct sunlight.

This datasheet offers general information. Please contact your local ECO representative for country-specific details.

Ensure the bag is closed securely after each use. Store in the original container.

General precautions

For animal treatment only. Keep out of reach and sight of children. Do not use after the expiry date stated on the label.

Package quantities

Aivlosin FG10 (8.5mg/g) is packed in 5 kg and 20kg bags.

Container disposal

Dispose of unused veterinary medicinal product or waste materials and empty packaging safely and in accordance with local waste disposal requirements.

Further information

It is recommended that Aivlosin is first mixed with 10kg of the feeding stuff, followed by the rest of the feeding stuff and mixed well. Medicated feed may be pelleted for up to 5 minutes at 70°C, which can be stored for 2 weeks to 3 months.

For further information contact:

ECO Animal Health
PO Box 47542
London N14 6WS
United Kingdom

Tel: +44 (0)208 447 8899

Fax: +44 (0)208 447 9292

Website: www.ecoanimalhealth.com and www.aivlosin.com or www.valosin.com